

Math 330 Homework 4

1. Let

$$A = \begin{bmatrix} -1 & 0 & 1 \\ 2 & 0 & -2 \\ 3 & 1 & 6 \end{bmatrix}, \quad x = \begin{bmatrix} -1 \\ 7 \\ -1 \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} 0 \\ 2 \\ 8 \end{bmatrix}.$$

(i) Find  $\frac{1}{3}x$

(ii) Find  $x + b$

(iii) Find  $x \cdot b$

(iv) Find  $\|b\|$

(v) Find  $Ax$

2. Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 1 & 3 & 1 \end{bmatrix}.$$

Write  $A$  as  $LDU$  where  $L$  is lower triangular with ones on its diagonal,  $D$  is diagonal and  $U$  is upper triangular with ones on its diagonal.

3. Let

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 & 2 \\ 1 & 2 & 3 & 4 & 5 \end{bmatrix}.$$

Find the reduced row echelon form  $R$  of  $A$ .

4. Consider the matrix  $A$  with reduced row echelon form  $R$  given by

$$A = \begin{bmatrix} 6 & 0 & 6 & 1 & 4 \\ 0 & 3 & 6 & 3 & 0 \\ 1 & 0 & 1 & 0 & 5 \\ 1 & 0 & 1 & 4 & 5 \end{bmatrix} \quad \text{and} \quad R = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

(i) Find a basis for the subspace  $\mathcal{C}(A)$  and state its dimension.

(ii) Find a basis for the subspace  $\mathcal{N}(A)$  and state its dimension.

5. Let

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 4 \end{bmatrix}.$$

Find  $\det A$ .

Math 330 Homework 4

6. Let

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}.$$

Find an orthogonal matrix  $Q$  and an upper triangular matrix  $R$  such that  $A = QR$ .

7. Let

$$Q = \begin{bmatrix} \frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ 0 & \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{2\sqrt{2}}{3} & \frac{-1}{3\sqrt{2}} & \frac{-1}{3\sqrt{2}} \end{bmatrix}, \quad R = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix}.$$

Note that  $Q$  is orthogonal and  $R$  upper triangular. Suppose  $A = QR$ . Find the  $x$  which minimizes  $\|Ax - b\|$ .

8. Let

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix}.$$

Find the eigenvectors and eigenvalues of  $A$

9. Let

$$A = \begin{bmatrix} 2 & 2 \\ -1 & 2 \end{bmatrix}.$$

Find the singular value decomposition  $A = U\Sigma V^T$  where  $U$  and  $V$  are orthogonal and  $\Sigma$  is diagonal.

10. Let  $u, v \in \mathbf{R}^2$  and  $\theta$  be the angle between  $u$  and  $v$ . Show that  $u \cdot v = \|u\|\|v\| \cos \theta$ .

11. Let  $A \in \mathbf{R}^{m \times n}$  and  $B \in \mathbf{R}^{l \times m}$ .

(i) Show that  $\mathcal{C}(BA) \subseteq \mathcal{C}(B)$ .

(ii) Given a concrete example where  $\mathcal{C}(B) \neq \mathcal{C}(BA)$ .

Math 330 Homework 4

**12.** Let  $A \in \mathbf{R}^{m \times n}$  where  $m \neq n$ . Suppose that the rank of  $A$  is  $\text{rank}(A) = r$ .

(i) What is  $\dim \mathcal{C}(A)$ ?

(ii) What is  $\dim \mathcal{N}(A)$ ?

(iii) What is  $\dim \mathcal{C}(A)^\perp$ ?

(iv) What is  $\dim \mathcal{N}(A)^\perp$ ?

(v) Show that  $\mathcal{C}(A)^\perp = \mathcal{N}(A^T)$ .

**13.** Let  $A \in \mathbf{R}^{4 \times 7}$ .

(i) Find the matrix  $E$  such that  $AE$  corresponds to the result obtained after performing the column operation  $c_2 \leftarrow c_2 - 3c_1$  on the matrix  $A$ .

(ii) Prove or disprove the claim that  $\mathcal{N}(AE) = \mathcal{N}(A)$ . If true explain why; if false provide a counterexample.