

Math 330: Final Exam Version A Sample Final

This is a closed-book closed-notes no-calculator-allowed in-class exam. Efforts have been made to keep the arithmetic simple. If it turns out to be complicated, that's either because I made a mistake or you did. In either case, do the best you can and check your work where possible. While getting the right answer is nice, this is not an arithmetic test. It's more important to clearly explain what you did and what you know.

1. Indicate in writing that you have understood the requirement to work independently by writing "I have worked independently on this quiz" followed by your signature as the answer to this question.

2. Write down the augmented matrix $[A|b]$ corresponding to the system of linear equations given by

from exam 1

$$\begin{cases} 2x_1 - 2x_2 + x_4 = 5 \\ x_2 - 5x_3 - 5x_4 = 2 \\ -2x_1 + 3x_2 - 5x_4 = -7 \end{cases}$$

but *do not* solve these equations.

use the rule $\det(AB) = (\det A)(\det B)$

3. Find $\det(A)$, $\det(B)$ and $\det(AB)$ where

from Exam 2

$$A = \begin{bmatrix} 1 & 2 & 17 \\ 0 & -2 & 8 \\ 0 & 0 & 3 \end{bmatrix}$$

and $B = \begin{bmatrix} 0 & 2 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

now what's $\det B =$

$$= \det \left(2 \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right)$$

corresponds to 3 row operations
 $r_1 \leftarrow 2r_1$
 $r_2 \leftarrow 2r_2$
 $r_3 \leftarrow 3r_3$

$$= 2^3 \det \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} = -8$$

Math 330: Final Exam Version A Sample Final

4. Consider the matrix A with reduced row echelon form R where

from Exam 2

$$A = \begin{bmatrix} \overset{P}{\frac{5}{2}} & \frac{4}{3} & -\frac{5}{2} & 0 & -\frac{2}{3} \\ -\frac{5}{4} & -\frac{2}{3} & \frac{5}{4} & \frac{4}{3} & -\frac{7}{6} \\ -\frac{15}{2} & -4 & \frac{15}{2} & \frac{4}{9} & \frac{1}{2} \end{bmatrix} \quad \text{and} \quad R = \begin{bmatrix} \overset{P}{1} & \overset{F}{\frac{8}{15}} & \overset{F}{-1} & \overset{P}{0} & \overset{P}{0} \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$x_1 \quad x_2 \quad x_3 \quad x_4 \quad x_5$

(i) Find a basis for $\text{Col}(A)$.

$$\left\{ \begin{bmatrix} 5/2 \\ -5/4 \\ -15/2 \end{bmatrix}, \begin{bmatrix} 0 \\ 4/3 \\ 4/9 \end{bmatrix}, \begin{bmatrix} -2/3 \\ -7/6 \\ 1/2 \end{bmatrix} \right\}$$

(ii) Find a basis for $\text{Nul}(A)$. $\approx \{x : Ax = 0\} \approx \{x : Rx = 0\}$

$$Rx = 0 \quad \begin{aligned} x_1 + \frac{8}{15}x_2 - x_3 &= 0 \\ x_4 &= 0 \\ x_5 &= 0 \end{aligned} \quad x = \begin{bmatrix} -\frac{8}{15}x_2 + x_3 \\ x_2 \\ x_3 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} -8/15 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} x_2 + \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} x_3$$

5. Let A be the matrix and x be the vector given by

New Problem

$$A = \begin{bmatrix} 2 & 6 & 7 \\ 3 & 2 & 7 \\ 5 & 6 & 4 \end{bmatrix} \quad \text{and} \quad x = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

Basis of $\text{Nul}(A)$: $\left\{ \begin{bmatrix} -8/15 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \right\}$

Show that x is an eigenvector of A and find the eigenvalue.

Math 330: Final Exam Version A Sample Final

6. Answer the following true false questions:

(i) Whenever a system has free variables, the solution set contains a unique solution.

(A) True

(B) False

(ii) An inconsistent system has more than one solution.

(A) True

(B) False

(iii) When two linear transformations are performed one after another, the combined effect may not always be a linear transformation.

(A) True

(B) False

(iv) $\det(A^{-1}) = 1/\det(A)$.

(A) True

(B) False

(v) Cramer's rule can only be used for invertible matrices.

(A) True

(B) False

(vi) If W is a subspace of \mathbf{R}^n and v is in both W and W^\perp , then $v = 0$.

(A) True

(B) False

(vii) If $A = QR$ where Q has orthonormal columns, then $R = Q^T A$.

(A) True

(B) False

(viii) If $A \in \mathbf{R}^{n \times n}$ is symmetric, there exists an orthonormal basis of \mathbf{R}^n which consists of eigenvectors of A .

(A) True

(B) False

(ix) Every matrix $A \in \mathbf{R}^{n \times n}$ can be factored as $A = SDS^{-1}$ where D is diagonal and S is an invertible matrix.

(A) True

(B) False

New

New

Math 330: Final Exam Version A Sample Final

7. Suppose $A \in \mathbf{R}^{2 \times 3}$ is given by

$$A = \begin{bmatrix} 1 & 3 & 2 \\ 3 & 5 & 1 \end{bmatrix}.$$

How many free variables does the equation $Ax = 0$ have? Find all solutions to the equation $Ax = 0$.

8. Suppose $A \in \mathbf{R}^{2 \times 2}$ is given by

$$A = \begin{bmatrix} 1 & 2 \\ -3 & 3 \end{bmatrix}.$$

Use the Gram-Schmidt algorithm to factor $A = QR$ where Q is a matrix with orthonormal columns and R is upper triangular.

9. Find the eigenvalues and eigenvectors of the matrix A where

$$A = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}.$$

Now

Solve $\det(A - \lambda I) = 0$ for λ then substitute those λ 's to find $\text{Nul}(A - \lambda I)$.

Math 330: Final Exam Version A Sample Final

10. The LU factorization of a matrix A is given by

$$L = \begin{bmatrix} 1 & 0 & 0 \\ -1/2 & 1 & 0 \\ 1/3 & -2/3 & 1 \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 5 & 6 \\ 0 & 0 & 7 \end{bmatrix}.$$

from
Exam 1

Explain how to use this factorization to solve the equation $Ax = b$ and then find the value of x corresponding to $b = (4, 6, 17)$.

Math 330: Final Exam Version A Sample Final

11. The QR factorization of a matrix A is given by

$$Q = \begin{bmatrix} \frac{1}{3} & \frac{2}{3\sqrt{5}} \\ -\frac{2}{3} & \frac{\sqrt{5}}{3} \\ \frac{2}{3} & \frac{4}{3\sqrt{5}} \end{bmatrix} \quad \text{and} \quad R = \begin{bmatrix} 3 & \frac{1}{3} \\ 0 & \frac{4\sqrt{5}}{3} \end{bmatrix}.$$

From Exam 2

I didn't clean up numbers, sorry...

Explain how to use this factorization to minimize $\|Ax - b\|$ and then find the minimizing value of x corresponding to $b = (1, 0, 1)$.

New

12. The matrix A given by

$$A = \begin{bmatrix} -1 & 4 & -2 \\ -3 & 4 & 0 \\ -3 & 1 & 3 \end{bmatrix}$$

has eigenvalues λ_i and eigenvectors x_i given by

$$\lambda_1 = 2, \quad x_1 = \begin{bmatrix} 2/3 \\ 1 \\ 1 \end{bmatrix}, \quad \lambda_2 = 3, \quad x_2 = \begin{bmatrix} 1/4 \\ 3/4 \\ 1 \end{bmatrix}, \quad \lambda_3 = 1, \quad x_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Find an invertible matrix S and a diagonal matrix D such that $A = SDS^{-1}$.

(i) What is D ?

(ii) What is S ?