

Math 330: Exam 2 Version C Sample Exam

This is a closed-book closed-notes no-calculator-allowed in-class exam. Efforts have been made to keep the arithmetic simple. If it turns out to be complicated, that's either because I made a mistake or you did. In either case, do the best you can and check your work where possible. While getting the right answer is nice, this is not an arithmetic test. It's more important to clearly explain what you did and what you know.

1. Indicate in writing that you have understood the requirement to work independently by writing "I have worked independently on this quiz" followed by your signature as the answer to this question.

2. Consider the matrix A with reduced row echelon form R where

$$A = \begin{bmatrix} 1 & 2 & -1 & -7 & 2 \\ -1 & -2 & 2 & 10 & -\frac{3}{2} \\ 2 & 4 & 1 & -5 & \frac{13}{2} \end{bmatrix} \quad \text{and} \quad R = \begin{bmatrix} 1 & 2 & 0 & -4 & 0 \\ 0 & 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

- (i) Find a basis for $\text{Col}(A)$.

- (ii) Find a basis for $\text{Nul}(A)$.

Math 330: Exam 2 Version C Sample Exam

3. Answer the following true false questions:

(i) $\det(A^{-1}) = (-1) \det(A)$.

(A) True

(B) False

(ii) Let S be an invertible matrix such that $AS = SB$. Then $\det A = \det B$.

(A) True

(B) False

(iii) If 0 is an eigenvalue of A , then A is invertible.

(A) True

(B) False

(iv) If A is invertible then A^T is also invertible.

(A) True

(B) False

4. Let λ be an eigenvalue of an invertible matrix A . Show λ^{-1} is an eigenvalue of A^{-1} .

5. How many pivot columns must a 5×7 matrix have if its columns span \mathbf{R}^5 ? Why?

Math 330: Exam 2 Version C Sample Exam

6. Find $\det(A)$, $\det(B)$ and $\det(AB)$ where

$$A = \begin{bmatrix} 2 & 1 & 6 \\ 0 & 3 & 5 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}.$$

7. The matrix A given by

$$A = \begin{bmatrix} 9 & -32 & -7 \\ 5 & -22 & -5 \\ -14 & 68 & 16 \end{bmatrix}$$

has eigenvalues λ_i and eigenvectors x_i given by

$$\lambda_1 = 2, \quad x_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \lambda_2 = 3, \quad x_2 = \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}, \quad \lambda_3 = -2, \quad x_3 = \begin{bmatrix} 1 \\ 1 \\ -3 \end{bmatrix}.$$

Find an invertible matrix S and a diagonal matrix D such that $A = SDS^{-1}$.

(i) What is D ?

(ii) What is S ?

Math 330: Exam 2 Version C Sample Exam

8. Find the eigenvalues and eigenvectors of the matrix A where

$$A = \begin{bmatrix} 7 & 4 \\ -8 & -5 \end{bmatrix}.$$